

Infrared regulators and SCET_{II}

Christian W. Bauer,^{*} Matthew P. Dorsten,[†] and Michael P. Salem[‡]
California Institute of Technology, Pasadena, CA 91125

We consider matching from SCET_I, which includes ultrasoft and collinear particles, onto SCET_{II} with soft and collinear particles at one loop. Keeping the external fermions off their mass shell does not regulate all IR divergences in both theories. We give a new prescription to regulate infrared divergences in SCET. Using this regulator, we show that soft and collinear modes in SCET_{II} are sufficient to reproduce all the infrared divergences of SCET_I. We explain the relationship between IR regulators and an additional mode proposed for SCET_{II}.

I. INTRODUCTION

Soft-collinear effective theory [1, 2, 3, 4] describes the interactions of soft and ultrasoft (usoft) particles with collinear particles. Using light-cone coordinates in which a general four-momentum is written as $p^\mu = (p^+, p^-, p^\perp) = (n \cdot p, \bar{n} \cdot p, p^\perp)$, where n and \bar{n} are four-vectors on the light cone ($n^2 = \bar{n}^2 = 0, n \cdot \bar{n} = 2$), these three degrees of freedom are distinguished by the scaling of their momenta:

$$\begin{aligned} \text{collinear: } p_c^\mu &\sim Q(\lambda^2, 1, \lambda), \\ \text{soft: } p_s^\mu &\sim Q(\lambda, \lambda, \lambda), \\ \text{usoft: } p_{us}^\mu &\sim Q(\lambda^2, \lambda^2, \lambda^2). \end{aligned} \tag{1}$$

The size of the expansion parameter λ is determined by the typical off-shellness of the collinear particles in a given problem. For example, in inclusive decays one typically has $p_c^2 \sim Q^2 \lambda^2 \sim Q \Lambda_{\text{QCD}}$, from which it follows that $\lambda = \sqrt{\Lambda_{\text{QCD}}/Q}$. For exclusive decays, however, one needs collinear particles with $p_c^2 \sim \Lambda_{\text{QCD}}^2$, giving $\lambda = \Lambda_{\text{QCD}}/Q$. One is usually interested in describing the interactions of these collinear degrees of freedom with non-perturbative degrees of freedom at rest, which satisfy $p^\mu \sim (\Lambda_{\text{QCD}}, \Lambda_{\text{QCD}}, \Lambda_{\text{QCD}})$. Thus inclusive processes involve interactions of collinear and usoft degrees of freedom, while exclusive decays are described by interactions of collinear and soft degrees of freedom. The theory describing the former set of degrees of freedom is called SCET_I, while the latter theory is called SCET_{II} [5].

Interactions between usoft and collinear degrees of freedom are contained in the leading-order Lagrangian of SCET_I,

$$\mathcal{L}_I = \bar{\xi}_n \left[in \cdot D + i \cancel{D}_c^\perp \frac{1}{i\bar{n} \cdot D_c} i \cancel{D}_c^\perp \right] \frac{\cancel{n}}{2} \xi_n, \tag{2}$$

and are well understood. The only interaction between collinear fermions and usoft gluons is from the derivative

$$iD^\mu = iD_c^\mu + gA_{us}^\mu. \tag{3}$$

These interactions can be removed from the leading-order Lagrangian by the field redefinition [4]

$$\begin{aligned} \xi_n &= Y_n \xi_n^{(0)}, \quad A_n = Y_n A_n^{(0)} Y_n^\dagger, \\ Y_n(x) &= \text{P exp} \left(ig \int_{-\infty}^0 ds n \cdot A_{us}(x + ns) \right). \end{aligned} \tag{4}$$

However, the same field redefinition has to be performed on the external operators in a given problem, and this reproduces the interactions with the usoft degrees of freedom. Consider for example the heavy-light current, which in SCET_I is given by

$$J_{hl}^I(\omega) = [\bar{\xi}_n W_n]_\omega \Gamma h_v, \tag{5}$$

^{*}Electronic address: bauer@theory.caltech.edu

[†]Electronic address: dorsten@theory.caltech.edu

[‡]Electronic address: salem@theory.caltech.edu

where h_v is the standard field of heavy quark effective theory [6], the Wilson line W_n is required to ensure collinear gauge invariance [3] and ω is the large momentum label of the gauge invariant $[\bar{\xi}_n W_n]$ collinear system. Written in terms of the redefined fields, this current is

$$J_{hl}^I(\omega) = \left[\bar{\xi}_n^{(0)} W_n^{(0)} \right]_{\omega} \Gamma [Y_n^\dagger h_v]. \quad (6)$$

For exclusive decays, we need to describe the interactions of soft with collinear particles. This theory is called SCET_{II} [5]. Since adding a soft momentum to a collinear particle takes this particle off its mass shell $(p_c + p_s)^2 \sim (Q\lambda, Q, Q\lambda)^2 \sim Q^2\lambda \sim Q\Lambda_{\text{QCD}}$, there are no couplings of soft to collinear particles in the leading-order Lagrangian.¹ Thus, the Lagrangian is given by [8, 9]

$$\mathcal{L}_{\text{II}} = \bar{\xi}_n \left[i n \cdot D_c + i \mathcal{D}_c^\perp \frac{1}{i \bar{n} \cdot D_c} i \mathcal{D}_c^\perp \right] \frac{\not{n}}{2} \xi_n. \quad (7)$$

In this theory, the heavy-light current is given by

$$J_{hl}^{II}(\omega, \kappa) = \left[\bar{\xi}_n^{(0)} W_n^{(0)} \right]_{\omega} \Gamma [S_n^\dagger h_v]_{\kappa}, \quad (8)$$

where S_n is a soft Wilson line in the n direction defined by

$$S_n(x) = \text{P exp} \left(ig \int_{-\infty}^0 ds n \cdot A_s(x + ns) \right). \quad (9)$$

This is the most general current invariant under collinear and soft gauge transformations.

This paper is organized as follows: We first consider matching the heavy-light current in SCET_I onto the heavy-light current in SCET_{II} using off-shell fermions. While the terms logarithmic in the off-shellness do not agree in the two theories, we argue that this is due to unregulated IR divergences in SCET_{II}. We then discuss IR regulators in SCET in more detail. We first identify the problems with SCET regulators and then propose a new regulator that addresses these issues. Using this regulator we then show that soft and collinear modes in SCET_{II} are sufficient to reproduce the IR divergences of SCET_I and explain the relationship between IR regulators and an additional mode proposed for SCET_{II} [9].

II. MATCHING FROM SCET_I ONTO SCET_{II}

The only difference between SCET_I and SCET_{II} is the typical off-shellness of the collinear degrees of freedom in the theory. The theory SCET_I allows fluctuations around the classical momentum with $p_c^2 \sim Q\Lambda_{\text{QCD}}$, while the theory SCET_{II} allows fluctuations with only $p_c^2 \sim \Lambda_{\text{QCD}}^2$. Since both theories expand around the same limit, SCET_{II} can be viewed as a low energy effective theory of SCET_I. Therefore, one can match from the theory SCET_I onto SCET_{II} by integrating out the $\mathcal{O}(\sqrt{Q\Lambda_{\text{QCD}}})$ fluctuations.

To illustrate this matching, we consider the heavy-light current. Using the definitions of this current given in Eqs. (5) and (8), we can write

$$J_{hl}^I(\omega) = \int d\kappa C(\omega, \kappa) J_{hl}^{II}(\omega, \kappa). \quad (10)$$

At tree level one finds trivially

$$C(\omega, \kappa) = 1. \quad (11)$$

In fact, this Wilson coefficient remains unity to all orders in perturbation theory, as was argued in Ref. [7].

To determine the matching coefficient at one loop, we calculate matrix elements of the current in the two theories. There are two diagrams in SCET_I, shown in Fig. 1. For on-shell external states, we find for the two integrals

$$iA^{Ia} = g^2 C_F \mu^{4-d} \int \frac{d^d k}{(2\pi)^d} \frac{1}{[-n \cdot k + i0][v \cdot k + i0][k^2 + i0]}, \quad (12)$$

$$iA^{Ib} = 2g^2 C_F \mu^{4-d} \int \frac{d^d k}{(2\pi)^d} \frac{\bar{n} \cdot (p_c - k)}{[-\bar{n} \cdot k + i0][k^2 - 2p_c \cdot k + i0][k^2 + i0]}. \quad (13)$$

¹ At higher orders, higher dimensional operators with at least two soft and two collinear particles can appear.



FIG. 1: Diagrams in SCET_I contributing to the matching. The solid square denotes an insertion of the heavy-light current.



FIG. 2: Diagrams in SCET_{II} contributing to the matching.

The diagrams in SCET_{II} are shown in Fig. 2. For on-shell external states the two integrals are exactly the same as in SCET_I :

$$iA^{IIa} = g^2 C_F \mu^{4-d} \int \frac{d^d k}{(2\pi)^d} \frac{1}{[-n \cdot k + i0][v \cdot k + i0][k^2 + i0]}, \quad (14)$$

$$iA^{IIb} = 2g^2 C_F \mu^{4-d} \int \frac{d^d k}{(2\pi)^d} \frac{\bar{n} \cdot (p_c - k)}{[-\bar{n} \cdot k + i0][k^2 - 2p_c \cdot k + i0][k^2 + i0]}. \quad (15)$$

Since the integrands are exactly the same, the loop diagrams will precisely cancel in the matching calculation. Thus we find that the Wilson coefficient $C(\omega, \kappa)$ remains unity, even at one loop. This confirms the arguments in [7] to this order.

The fact that both of these integrals are scaleless and therefore zero might bother some readers. The vanishing of these diagrams is due to the cancellation of collinear, infrared (IR) and ultraviolet (UV) divergences. Introducing an IR regulator will separate these divergences, and the UV will be regulated by dimensional regularization. In Ref. [1] a small off-shellness was introduced to regulate the IR divergences of SCET_I . In Refs. [9] the divergence structure of SCET_{II} was studied keeping both the heavy and the collinear fermions off-shell. Using this IR regulator, the authors of Refs. [9] argued that SCET_{II} does not reproduce the IR divergences of SCET_I and introduced a new mode in SCET_{II} to fix this problem. To gain more insight into their argument, we will go through their calculation in some detail.

In SCET_I the first diagram is

$$\begin{aligned} A_{p_c}^{Ia} &= -ig^2 C_F \mu^{4-d} \int \frac{d^d k}{(2\pi)^d} \frac{1}{[\tilde{p}_c - n \cdot k + i0][v \cdot (p_s - k) + i0][k^2 + i0]} \\ &= -\frac{g^2 C_F}{2\pi} (4\pi)^{1-d/2} \Gamma\left(2 - \frac{d}{2}\right) \mu^{4-d} \int_0^\infty dn \cdot k (n \cdot k - \tilde{p}_c)^{-1} n \cdot k^{d/2-2} (n \cdot k - 2v \cdot p_s)^{d/2-2} \\ &= \frac{\alpha_s C_F}{4\pi} \left[-\frac{1}{\epsilon^2} + \frac{2}{\epsilon} \log \frac{-\tilde{p}_c}{\mu} - 2 \log^2 \frac{-\tilde{p}_c}{\mu} + 2 \log \left(1 - \frac{2v \cdot p_s}{\tilde{p}_c}\right) \log \frac{2v \cdot p_s}{\tilde{p}_c} + 2 \text{Li}_2 \left(\frac{2v \cdot p_s}{\tilde{p}_c}\right) - \frac{3\pi^2}{4} \right], \end{aligned} \quad (16)$$

where $d = 4 - 2\epsilon$ and

$$\tilde{p}_c = \frac{p_c^2}{\bar{n} \cdot p_c}. \quad (17)$$

In going from the first line to the second, we closed the $\bar{n} \cdot k$ contour below, thus restricting $n \cdot k$ to positive values, and

performed the Euclidean k_\perp integral. The second diagram gives

$$\begin{aligned} A_{p_c}^{Ib} &= -2ig^2 C_F \mu^{4-d} \int \frac{d^d k}{(2\pi)^d} \frac{\bar{n} \cdot (p_c - k)}{[-\bar{n} \cdot k + i0][(k - p_c)^2 + i0][k^2 + i0]} \\ &= \frac{\alpha_s C_F}{4\pi} \left[\frac{2}{\epsilon^2} + \frac{2}{\epsilon} - \frac{2}{\epsilon} \log \frac{-p_c^2}{\mu^2} + \log^2 \frac{-p_c^2}{\mu^2} - 2 \log \frac{-p_c^2}{\mu^2} + 4 - \frac{\pi^2}{6} \right]. \end{aligned} \quad (18)$$

In this diagram it is necessary to choose $d < 4$ for the k_\perp integral, but one requires $d > 4$ for the $\bar{n} \cdot k$ integral. In the former integral, dimensional regularization regulates the divergence at $k_\perp = \infty$, while in the latter it regulates the divergence at $\bar{n} \cdot k = 0$. Both of these divergences have to be interpreted as UV, as discussed in section III. Each diagrams contain a mixed UV-IR divergence of the form $\log p_c^2/\epsilon$. This mixed divergence cancels in the sum of the two diagrams and we find, after also adding the wave function contributions,

$$\begin{aligned} A_{p_c}^I &= \frac{\alpha_s C_F}{4\pi} \left[\frac{1}{\epsilon^2} + \frac{2}{\epsilon} \log \frac{\mu}{\bar{n} \cdot p_c} + \frac{5}{2\epsilon} + \log^2 \frac{-p_c^2}{\mu^2} - 2 \log^2 \frac{-\tilde{p}_c}{\mu} - \frac{3}{2} \log \frac{-p_c^2}{\mu^2} - 2 \log \frac{-2v \cdot p_s}{\mu} \right. \\ &\quad \left. + 2 \log \left(1 - \frac{2v \cdot p_s}{\tilde{p}_c} \right) \log \frac{2v \cdot p_s}{\tilde{p}_c} + 2 \text{Li}_2 \left(\frac{2v \cdot p_s}{\tilde{p}_c} \right) + \frac{11}{2} - \frac{11\pi^2}{12} \right]. \end{aligned} \quad (19)$$

This reproduces the IR behavior of full QCD.

Now consider the SCET_{II} diagrams. The second is identical to the one in SCET_I :

$$A_{p_c}^{IIb} = A_{p_c}^{Ib}. \quad (20)$$

The first diagram, however, is different and we find

$$\begin{aligned} A_{p_c}^{IIa} &= -ig^2 C_F \mu^{4-d} \int \frac{d^d k}{(2\pi)^d} \frac{1}{[-n \cdot k + i0][v \cdot (p_s - k) + i0][k^2 + i0]} \\ &= -\frac{g^2 C_F}{2\pi} \mu^{4-d} \int_0^\infty dn \cdot k \int \frac{d^{d-2} k_\perp}{(2\pi)^{d-2}} \frac{1}{n \cdot k (k_\perp^2 + n \cdot k^2 - 2v \cdot p_s n \cdot k)} \\ &= -\frac{\alpha_s C_F}{2\pi} (4\pi)^{2-d/2} \Gamma \left(2 - \frac{d}{2} \right) \mu^{4-d} \int_0^\infty dn \cdot k n \cdot k^{d/2-3} (n \cdot k - 2v \cdot p_s)^{d/2-2}. \end{aligned} \quad (21)$$

Note that convergence of this integral at $n \cdot k = \infty$ requires $d < 4$, whereas convergence at $n \cdot k = 0$ requires $d > 4$. In this case dimensional regularization is regulating both a UV divergence at $n \cdot k = \infty$, as well as the divergence at $n \cdot k = 0$, which is IR in nature, as we will discuss in section III. Using the variable transformation $x = n \cdot k / (n \cdot k - 2v \cdot p_s)$ to relate this integral to a beta function [10] one finds

$$A_{p_c}^{IIa} = \frac{\alpha_s C_F}{4\pi} \left[\frac{1}{\epsilon^2} - \frac{2}{\epsilon} \log \frac{-2v \cdot p_s}{\mu} + 2 \log^2 \frac{-2v \cdot p_s}{\mu} + \frac{5\pi^2}{12} \right]. \quad (22)$$

Adding the two diagrams together with the wave function contributions gives

$$\frac{\alpha_s C_F}{4\pi} \left[\frac{3}{\epsilon^2} - \frac{2}{\epsilon} \log \frac{2v \cdot p_s p_c^2}{\mu^3} + \frac{5}{2\epsilon} + \log^2 \frac{-p_c^2}{\mu^2} + 2 \log^2 \frac{-2v \cdot p_s}{\mu} - \frac{3}{2} \log \frac{-p_c^2}{\mu^2} - 2 \log \frac{-2v \cdot p_s}{\mu} + \frac{11}{2} + \frac{\pi^2}{4} \right]. \quad (23)$$

We can see that in the sum of the two diagrams the terms proportional to $\log p_c^2/\epsilon$ or $\log v \cdot p_s/\epsilon$ do not cancel as they did in SCET_I. Furthermore, the finite terms logarithmic in p_c^2 or $v \cdot p_s$ do not agree with the corresponding terms in




FIG. 3: Contribution of the additional SCET_{II} mode proposed in Refs. [9].

the SCET_I result. This fact prompted the authors of Refs. [9] to conclude that SCET_{II} does not reproduce the IR divergences of SCET_{II} and that a new mode is needed in the latter effective theory. However, as we mentioned above, there are problems with IR effects in this diagram. In fact, as we will show in great detail in the next section, the off-shellness of the fermions does not regulate all IR divergences in this diagram. This means that the fact that the terms logarithmic in the fermion off-shellness do not agree between SCET_I and SCET_{II} does not imply that the IR divergences are not reproduced correctly since some $1/\epsilon$ poles are IR in origin.

We also calculate the diagram in SCET_{II} containing the additional mode proposed in Refs. [9]. The new messenger mode has momenta scaling as $p_{sc}^\mu \sim (\Lambda_{\text{QCD}}^2/Q, \Lambda_{\text{QCD}}, \Lambda_{\text{QCD}}^{3/2}/Q^{1/2})$. (Note that the invariant mass of this term satisfies $p_{sc}^2 \ll \Lambda_{\text{QCD}}^2$.) The diagram is shown in Fig. 3 and we find

$$\begin{aligned} A_{p_c}^{IIc} &= -2ig^2 C_F \mu^{4-d} \int \frac{d^d k}{(2\pi)^d} \frac{1}{[\tilde{p}_c - n \cdot k + i0][2v \cdot p_s - \bar{n} \cdot k + i0][k^2 + i0]} \\ &= \frac{\alpha_s C_F}{4\pi} \left[-\frac{2}{\epsilon^2} + \frac{2}{\epsilon} \log \frac{2v \cdot p_s \tilde{p}_c}{\mu^2} - \log^2 \frac{2v \cdot p_s \tilde{p}_c}{\mu^2} - \frac{\pi^2}{2} \right]. \end{aligned} \quad (24)$$

Adding this term to the Eq. (23) cancels the terms proportional to $\log(2v \cdot p_s p_c^2/\mu^3)/\epsilon$ and we find

$$\begin{aligned} A_{p_c}^{II} &= \frac{\alpha_s C_F}{4\pi} \left[\frac{1}{\epsilon^2} + \frac{2}{\epsilon} \log \frac{\mu}{\bar{n} \cdot p_c} + \frac{5}{2\epsilon} + \log^2 \frac{-p_c^2}{\mu^2} - \frac{3}{2} \log \frac{-p_c^2}{\mu^2} - 2 \log \frac{-2v \cdot p_s}{\mu} \right. \\ &\quad \left. + 2 \log^2 \left(\frac{-2v \cdot p_s}{\mu} \right) - \log^2 \left(\frac{2v \cdot p_s \tilde{p}_c}{\mu^2} \right) + \frac{11}{2} - \frac{\pi^2}{4} \right]. \end{aligned} \quad (25)$$

This result does not agree with the SCET_I expression in Eq. (19). However, this is expected, since the off-shellness in SCET_{II} satisfies $\tilde{p}_c \ll v \cdot p_s$. In this limit the SCET_I result in Eq. (19) agrees with the result in Eq. (25).

III. INFRARED REGULATORS IN SCET

A. Problems with known IR regulators

One of the most important properties of SCET_I is the field redefinition given in Eq. (4), which decouples the usoft from the collinear fermions. It is the crucial ingredient for proving factorization theorems. Furthermore, only after performing this field redefinition is it simple to match from SCET_I onto SCET_{II}, since one can identify the Wilson line Y_n in SCET_I with the Wilson line S_n in SCET_{II}. However, it is a well known fact that field redefinitions only leave on-shell Green functions invariant [11]. Hence, the off-shellness of the collinear quark p_c^2 used to regulate the IR in SCET_I takes away our ability to perform this field redefinition. Since no field redefinition is performed on the heavy quark, one is free to give it an off-shellness.

IR divergences appear in individual diagrams, but they cancel in the set of diagrams contributing to a physical observable. More specifically, the IR divergences in virtual loop diagrams are cancelled against those in real emissions, which physically have to be included due to finite detector resolutions. From this it is obvious that the IR divergences in the heavy-light current originate from regions of phase space where either the gluon three-momentum $|\mathbf{k}|$ or the angle θ between the gluon and the light fermion goes to zero. Other divergences arise if the three-momentum of the gluon goes to infinity or θ goes to π . These divergences are UV. To check if the IR divergences match between the two theories one has to use an IR regulator that regulates all IR divergences in both theories. To get insight into the behavior of the three-momentum and the angle, it will be instructive to perform the required loop integrals by integrating over k_0 using the method of residues, and then integrating over the magnitude of the three-momentum and the solid angle. This will allow us to identify clearly the IR divergences as described above. Let us illustrate this method by showing that all $1/\epsilon$ divergences in the SCET_I one-loop calculation of the previous section are UV. For the first diagram we find

$$\begin{aligned} A_{p_c}^{Ia} &= -ig^2 C_F \mu^{4-d} \int \frac{d^d k}{(2\pi)^d} \frac{1}{[\tilde{p}_c - n \cdot k + i0][v \cdot (p_s - k) + i0][k^2 + i0]} \\ &= -\frac{g^2 C_F}{2} \frac{\Omega_{d-2}}{(2\pi)^{d-1}} \mu^{4-d} \int_0^\infty d|\mathbf{k}| |\mathbf{k}|^{d-2} \int_{-1}^1 d\cos\theta \sin^{d-4}\theta \frac{1}{(|\mathbf{k}|(1 - \cos\theta) - \tilde{p}_c)(|\mathbf{k}| - v \cdot p_s)|\mathbf{k}|}. \end{aligned} \quad (26)$$

Performing the remaining integrals, we of course reproduce the result obtained previously, but this form demonstrates that all divergences from regions $|\mathbf{k}| \rightarrow 0$ and $(1 - \cos\theta) \rightarrow 0$ are regulated by the infrared regulators and thus all $1/\epsilon$ divergences are truly UV.

The second diagram is

$$\begin{aligned} A_{p_c}^{Ib} &= -2ig^2 C_F \mu^{4-d} \int \frac{d^d k}{(2\pi)^d} \frac{\bar{n} \cdot (p_c - k)}{[-\bar{n} \cdot k + i0][(p_c - k)^2 + i0][k^2 + i0]} \\ &= -2ig^2 C_F [I_1 + I_2], \end{aligned} \quad (27)$$

where I_1 and I_2 are the integrals with the $\bar{n} \cdot p$ and the $\bar{n} \cdot k$ terms in the numerator, respectively. The integral I_2 is standard and we find

$$I_2 = \frac{i}{16\pi^2} \left[\frac{1}{\epsilon} - \log \frac{-p_c^2}{\mu^2} + 2 \right], \quad (28)$$

where ϵ regulates only UV divergences. For the first integral we again perform the k_0 integral by contours and we find

$$\begin{aligned} I_1 &= \frac{i\bar{n} \cdot p_c}{2} \frac{\Omega_{d-2}}{(2\pi)^{d-1}} \mu^{4-d} \int_0^\infty d|\mathbf{k}| |\mathbf{k}|^{d-2} \int_{-1}^1 d\cos\theta \sin^{d-4}\theta \\ &\quad \left[-\frac{1}{\mathbf{k}^2(1+\cos\theta)[2|\mathbf{k}|(p_0 - |\mathbf{p}|\cos\theta) - p_c^2]} + \frac{1}{a[p_0 + a + |\mathbf{k}|\cos\theta][2p_0^2 + 2p_0a - 2|\mathbf{k}||\mathbf{p}|\cos\theta - p_c^2]} \right], \end{aligned} \quad (29)$$

where

$$p_c = (p_0, \mathbf{p}), \quad a = \sqrt{\mathbf{k}^2 + \mathbf{p}^2 - 2|\mathbf{k}||\mathbf{p}|\cos\theta}. \quad (30)$$

From this expression we can again see that all IR singularities from $|\mathbf{k}| \rightarrow 0$ and $(1 - \cos\theta) \rightarrow 0$ are regulated by the off-shellness, and all remaining divergences are UV. Note furthermore that in the limit $|\mathbf{k}| \rightarrow \infty$, with unrestricted θ , the two terms cancel each other, so that there is no usual UV divergence. This agrees with the fact that there are five powers of k in the denominator of the integrand in Eq. (27). However, in the limit $|\mathbf{k}| \rightarrow \infty$ with $|\mathbf{k}|\cos\theta \rightarrow 0$ the second term of Eq. (29) remains finite, whereas the first term develops a double divergence. Thus, it is this region of phase space that gives rise to the double pole in this diagram. The presence of the square roots makes the evaluation of the remaining integrals difficult, but we have checked that we reproduce the divergent terms of the result given in Eq. (18).

From the above discussion it follows that the off-shellness of the external fermions regulates all the IR divergences, and that the $1/\epsilon$ divergences all correspond to divergences of UV origin. The situation is different in SCET_{II}, since the off-shellness of the light quark does not enter diagram (a). We find

$$\begin{aligned} A_{p_c}^{IIa} &= -ig^2 C_F \mu^{4-d} \int \frac{d^d k}{(2\pi)^d} \frac{1}{[-n \cdot k + i0][v \cdot (p_s - k) + i0][k^2 + i0]} \\ &= -\frac{g^2 C_F}{2} \frac{\Omega_{d-2}}{(2\pi)^{d-1}} \mu^{4-d} \int_0^\infty d|\mathbf{k}| |\mathbf{k}|^{d-2} \int_{-1}^1 d\cos\theta \sin^{d-4}\theta \frac{1}{\mathbf{k}^2(1-\cos\theta)(|\mathbf{k}| - v \cdot p_s)}. \end{aligned} \quad (31)$$

The IR divergence originating from the limit $(1 - \cos\theta) \rightarrow 0$ is not regulated by the off-shellness. Thus part of the $1/\epsilon$ divergences in Eq. (22) are of IR origin. In other words, the fact that the terms logarithmic in the off-shellness in the SCET_I amplitude Eq. (19) are not reproducing the corresponding terms in the SCET_{II} amplitude Eq. (23) does not imply that the IR divergences do not match between the two theories. In order to check whether the IR divergences of the two theories match, one needs a regulator that regulates all IR divergences in both SCET_I and SCET_{II}.

As an alternative IR regulator one could try to use a small gluon mass. Consider the first diagram in SCET_I again, this time with a gluon mass. We find

$$\begin{aligned} A_m^{Ia} &= -ig^2 C_F \mu^{4-d} \int \frac{d^d k}{(2\pi)^d} \frac{1}{[-n \cdot k + i0][v \cdot (p_s - k) + i0][k^2 - m^2 + i0]} \\ &= -\frac{g^2 C_F}{2} \frac{\Omega_{d-2}}{(2\pi)^{d-1}} \mu^{4-d} \int_0^\infty d|\mathbf{k}| |\mathbf{k}|^{d-2} \int_{-1}^1 d\cos\theta \sin^{d-4}\theta \frac{1}{(\mathbf{k}^2 + m^2 - v \cdot p_s \sqrt{\mathbf{k}^2 + m^2})(\sqrt{\mathbf{k}^2 + m^2} - |\mathbf{k}|\cos\theta)}. \end{aligned} \quad (32)$$

Again, all divergences $|\mathbf{k}| \rightarrow 0$ and $(1 - \cos\theta) \rightarrow 0$ are regulated by the gluon mass, but in the limit $|\mathbf{k}| \rightarrow \infty$ with $|\mathbf{k}|\cos\theta \rightarrow 0$ the integrand becomes

$$\frac{|\mathbf{k}|^{d-4} \sin^{d-4}\theta}{|\mathbf{k}|\cos\theta + \frac{m^2}{2|\mathbf{k}|}}, \quad (33)$$

so that the term that could potentially regulate the $(1 - \cos\theta) \rightarrow 0$ divergence goes to zero as $|\mathbf{k}| \rightarrow \infty$. This is why a gluon mass cannot be used to regulate the IR of SCET.

B. A new regulator for SCET

The gluon mass is not an appropriate IR regulator for SCET because it appears in the combination $m^2/|\mathbf{k}|$ in the expression (33). Instead of using a gluon mass, consider adding the terms ²

$$\begin{aligned}\mathcal{L}_{\text{reg}}^c &= -\frac{\delta}{2}A_\mu^c\bar{\mathcal{P}}A_c^\mu \\ \mathcal{L}_{\text{reg}}^{(u)s} &= -\frac{\delta}{2}A_\mu^{(u)s}i\bar{n}\cdot\partial A_{(u)s}^\mu\end{aligned}\quad (34)$$

to the collinear and (u)soft gluon Lagrangians. Here, $\bar{\mathcal{P}}$ is the label operator which picks out the large momentum label of the collinear gluon field. Both of these terms are generated if a similar term is added to the full QCD gluon action before constructing SCET. Note that these terms preserve the invariance of the theory under the field redefinitions given in Eq. (4).

The infinitesimal, dimensionful parameter δ suffices to regulate all IR divergences in SCET, unlike the gluon mass. Following the same steps as in Eq. (32). We find

$$A_\delta^{Ia} = -\frac{g^2C_F}{2}\frac{\Omega_{d-2}}{(2\pi)^{d-1}}\mu^{4-d}\int_0^\infty d|\mathbf{k}||\mathbf{k}|^{d-2}\int_{-1}^1 d\cos\theta\sin^{d-4}\theta\frac{8}{b(\delta+b)(\delta+b-2|\mathbf{k}|\cos\theta)},\quad (35)$$

where

$$b = \sqrt{4\mathbf{k}^2 + \delta^2 + 4|\mathbf{k}|\delta\cos\theta}.\quad (36)$$

Obviously, the parameter δ regulates the divergences $|\mathbf{k}| \rightarrow 0$ and $(1 - \cos\theta) \rightarrow 0$, just as the gluon mass did. Expanding around the limit $|\mathbf{k}| \rightarrow \infty$ with $|\mathbf{k}|\cos\theta \rightarrow 0$ the integrand becomes

$$\frac{|\mathbf{k}|^{d-4}\sin^{d-4}\theta}{|\mathbf{k}|\cos\theta + \delta},\quad (37)$$

and this IR region is therefore regulated as well. Even though δ is enough to regulate all IR divergences in SCET, we will keep the heavy quark off its mass-shell for later convenience.

Performing the integrals using the method above is difficult. While performing the k_0 integration using the method of residues gives insight into the divergence structure of the loop integrals, it is simpler to perform the integrals using the variables $n\cdot k$ and $\bar{n}\cdot k$ instead. The first diagram in SCET_I with this new regulator is then given by

$$\begin{aligned}A_\delta^{Ia} &= -ig^2C_F\mu^{4-d}\int\frac{d^dk}{(2\pi)^d}\frac{1}{[-n\cdot k+i0][v\cdot(p_s-k)+i0][k^2-\delta\bar{n}\cdot k+i0]} \\ &= -\frac{g^2C_F}{2\pi}(4\pi)^{1-d/2}\Gamma\left(2-\frac{d}{2}\right)\mu^{4-d}\int_\delta^\infty dn\cdot k\int_\delta^\infty d\bar{n}\cdot k\frac{1}{(n\cdot k-\delta)^{d/2-2}(n\cdot k-2v\cdot p_s)^{d/2-2}} \\ &= \frac{\alpha_s C_F}{4\pi}\left[-\frac{1}{\epsilon^2}+\frac{2}{\epsilon}\log\frac{\delta}{\mu}-2\log^2\frac{\delta}{\mu}+2\log\left(1-\frac{2v\cdot p_s}{\delta}\right)\log\frac{2v\cdot p_s}{\delta}+2\text{Li}_2\left(1-\frac{2v\cdot p_s}{\delta}\right)-\frac{3\pi^2}{4}\right].\end{aligned}\quad (38)$$

Similarly, it is possible to show that the parameter δ regulates all IR divergences in the second diagram, for which we find

$$\begin{aligned}A_\delta^{Ib} &= -2ig^2C_F\mu^{4-d}\int\frac{d^dk}{(2\pi)^d}\frac{\bar{n}\cdot(p_c-k)}{[-\bar{n}\cdot k+i0][k^2-2k\cdot p_c+i0][k^2-\delta\bar{n}\cdot k+i0]} \\ &= \frac{\alpha_s C_F}{4\pi}\left[\frac{2}{\epsilon^2}+\frac{2}{\epsilon}-\frac{2}{\epsilon}\log\frac{\delta\bar{n}\cdot p_c}{\mu^2}+\log^2\frac{\delta\bar{n}\cdot p_c}{\mu^2}-2\log\frac{\delta\bar{n}\cdot p_c}{\mu^2}+4-\frac{\pi^2}{6}\right].\end{aligned}\quad (39)$$

² An alternative regulator has been introduced in Ref. [12]. In that paper a quark mass is used in conjunction with an ‘‘analytic’’ regulator, which regulates the $(1 - \cos\theta) \rightarrow 0$ divergence. The conclusions about the soft-collinear mode in Ref. [9] are similar to the ones drawn here. However, we believe that a regulator such as the one introduced here is advantageous, since it can naturally be defined at the level of the Lagrangian, and a single dimensionful parameter regulates all IR divergences.

The mixed UV-IR divergences cancel in the sum of the two diagrams,

$$A_\delta^I = \frac{\alpha_s C_F}{4\pi} \left[\frac{1}{\epsilon^2} + \frac{2}{\epsilon} \log \frac{\mu}{\bar{n} \cdot p_c} + \frac{5}{2\epsilon} + \log^2 \frac{\delta \bar{n} \cdot p_c}{\mu^2} - 2 \log^2 \frac{\delta}{\mu} - \frac{3}{2} \log \frac{\delta \bar{n} \cdot p_c}{\mu^2} - 2 \log \frac{\delta - 2v \cdot p_s}{\mu} \right. \\ \left. + 2 \log \left(1 - \frac{2v \cdot p_s}{\delta} \right) \log \frac{2v \cdot p_s}{\delta} + 2 \text{Li}_2 \left(1 - \frac{2v \cdot p_s}{\delta} \right) + \frac{11}{2} - \frac{11\pi^2}{12} \right], \quad (40)$$

and one can show that this result reproduces the IR behavior of full QCD.

Since the regulator is in the gluon action, it is the same for SCET_I and SCET_{II}, and the two diagrams in SCET_{II} are identical to those in SCET_I since the integrands are exactly equal:

$$A_\delta^{II} = A_\delta^I. \quad (41)$$

Therefore, the IR divergences in SCET_{II} are exactly the same as those in SCET_I.

While in SCET_I it is possible to choose the scaling $\delta \sim Q\lambda^2$ such that both the contributions to the collinear and the usoft gluon action are leading order in the power counting, the same is not true in SCET_{II}. Choosing $\delta \sim Q\lambda^2$ to make the IR regulator leading order in collinear gluon Lagrangian makes it suppressed by one power of λ in the soft Lagrangian. This can be understood physically, since in going from SCET_I to SCET_{II} the typical scaling of the (u)soft momenta remains of order Λ_{QCD} , while the off-shellness of the collinear particles is lowered. However, the IR divergence from $n \cdot k \rightarrow 0$ corresponds to $(1 - \cos \theta) \rightarrow 0$, and the typical cutoff on $(1 - \cos \theta)$ is set by the collinear scales. Since $n \cdot k_c \ll n \cdot k_s$ it is natural that any cutoff κ regulating the $n \cdot k_s \rightarrow 0$ divergence will satisfy $\kappa \ll n \cdot k_s$. This is not a problem, since the IR regulator does not introduce a new scale into the effective theory.

If one insists on keeping the scaling manifest, one is forced to drop the regulator term in the soft gluon Lagrangian. In this case, the diagram (a) in SCET_{II} no longer includes the IR regulator δ and is therefore not regulated properly. The calculation then reduces to the result given in Eq. (22). Part of the $1/\epsilon$ divergences in this result are from true UV divergences, but others are due to the unregulated $(1 - \cos \theta) \rightarrow 0$ IR divergences, which arise from physics at the scale $n \cdot k \sim Q\lambda^2$. These IR divergences can be recovered by adding a diagram containing a gluon scaling as $n \cdot k \sim Q\lambda^2$, $\bar{n} \cdot k \sim Q\lambda$. Requiring $n \cdot k \bar{n} \cdot k \sim k_\perp^2$, this is the soft collinear messenger mode introduced in [9]. The resulting diagram (c) gives

$$A_\delta^{IIc} = -2ig^2 C_F \mu^{4-d} \int \frac{d^d k}{(2\pi)^d} \frac{1}{[-n \cdot k + i0][2v \cdot p_s - \bar{n} \cdot k + i0][k^2 - \delta \bar{n} \cdot k + i0]} \\ = \frac{\alpha_s C_F}{4\pi} \left[-\frac{2}{\epsilon^2} + \frac{2}{\epsilon} \log \frac{-2v \cdot p_s \delta}{\mu^2} - \log^2 \frac{-2v \cdot p_s \delta}{\mu^2} - \frac{\pi^2}{2} \right]. \quad (42)$$

Adding all the diagrams we find

$$\frac{\alpha_s C_F}{4\pi} \left[\frac{1}{\epsilon^2} + \frac{2}{\epsilon} \log \frac{\mu}{\bar{n} \cdot p_c} + \frac{5}{2\epsilon} + \log^2 \frac{\delta \bar{n} \cdot p_c}{\mu^2} - \frac{3}{2} \log \frac{\delta \bar{n} \cdot p_c}{\mu^2} - 2 \log \frac{-2v \cdot p_s}{\mu} \right. \\ \left. + 2 \log^2 \left(\frac{-2v \cdot p_s}{\mu} \right) - \log^2 \left(\frac{-2v \cdot p_s \delta}{\mu^2} \right) + \frac{11}{2} - \frac{\pi^2}{4} \right], \quad (43)$$

which again reproduces the SCET_I result for $\delta \ll v \cdot p_s$. From this discussion it follows that the presence of the soft collinear messenger mode depends on the precise implementation of the IR regulator in the theory. Since the definition of an effective theory should be independent of the regulator used for an explicit calculation, one can view the soft-collinear messenger mode as part of the IR regulator.

The term added to the gluon Lagrangian breaks gauge invariance. However, in this regard it is on the same footing as a gluon mass. Since the coupling of gluons to fermions is via a conserved current, this breaking of gauge invariance is only a problem once gluon self-interactions are taken into account. For the renormalization of operators such as the heavy-light current considered in this paper, this only arises at the two loop level. In matching calculations, the IR divergences always cancel. Hence any IR regulator, including the one proposed here is applicable to matching calculations at any order in perturbation theory. More care has to be taken when using this regulator to calculate operator mixing, and in this case gauge non-invariant operators have to be included beyond one loop order. The main advantage of the new regulator is that it preserves invariance of SCET_I under the field redefinition given in Eq. (4).

IV. CONCLUSIONS

We have considered the matching of the heavy-light current in SCET_I onto the corresponding current in SCET_{II}, in particular addressing the question whether all long distance physics in SCET_I is correctly reproduced in SCET_{II}.

Using the off-shellness of the external heavy and light fermions, it was argued in Refs. [9] that a new collinear-soft messenger mode is required in SCET_{II} to reproduce all the long distance physics of SCET_I. Regulating the IR in SCET_{II} with an off-shellness is problematic, since the off-shellness prevents performing the field redefinition required to decouple the usoft gluons from the collinear particles, which allows the matching onto SCET_{II} easily. In this paper we investigated the relationship between IR regulators and the definition of SCET_{II}. By performing the k_0 loop integral by contours and then writing the remaining integrals as $d|\mathbf{k}|d\cos\theta$, we showed explicitly that the off-shellness leaves the IR angular divergence $(1 - \cos\theta) \rightarrow 0$ unregulated in SCET_{II}.

We then introduced a new regulator for SCET that regulates soft ($|\mathbf{k}| \rightarrow 0$) and collinear ($\cos\theta \rightarrow 1$) IR divergences in both SCET_I and SCET_{II}. This regulator modifies the gluon action, much like a gluon mass, and thus preserves the field redefinition required to decouple usoft gluons from collinear particles in SCET. Using this regulator, we showed explicitly that SCET_{II} as formulated in Refs. [4, 7] reproduces all the IR divergences of SCET_I. We also argued that any cutoff κ regulating the collinear divergence has to satisfy $\kappa \ll n \cdot k_s$. Regulating SCET_{II} this way therefore naturally requires keeping a formally subleading regulator in the theory.

We also showed that a soft-collinear messenger mode is required in the definition of the IR regulator if one insists on power counting the regulator in the same way as kinetic terms in the action. In this case, there are unregulated IR divergences left in soft diagrams, which are corrected by additional contributions from the soft-collinear mode.

The new regulator introduced in this paper preserves the invariance of SCET_I under the field redefinitions (4), and is therefore useful in studying factorization theorems beyond tree level.

Acknowledgments

We would like to thank Thomas Becher, Richard Hill, Bjorn Lange, Michael Luke, Matthias Neubert, David Politzer, Ira Rothstein, Iain Stewart and Mark Wise for useful discussions. This work was supported by the Department of Energy under the contract DE-FG03-92ER40701.

-
- [1] C. W. Bauer, S. Fleming and M. E. Luke, Phys. Rev. D **63**, 014006 (2001).
 - [2] C. W. Bauer, S. Fleming, D. Pirjol and I. W. Stewart, Phys. Rev. D **63**, 114020 (2001).
 - [3] C. W. Bauer and I. W. Stewart, Phys. Lett. B **516**, 134 (2001).
 - [4] C. W. Bauer, D. Pirjol and I. W. Stewart, Phys. Rev. D **65**, 054022 (2002).
 - [5] C. W. Bauer, D. Pirjol and I. W. Stewart, Phys. Rev. D **67**, 071502 (2003) [arXiv:hep-ph/0211069].
 - [6] A. V. Manohar and M. B. Wise, Cambridge Monogr. Part. Phys. Nucl. Phys. Cosmol. **10**, 1 (2000).
 - [7] C. W. Bauer, D. Pirjol and I. W. Stewart, Phys. Rev. D **68**, 034021 (2003).
 - [8] R. J. Hill and M. Neubert, Nucl. Phys. B **657**, 229 (2003) [arXiv:hep-ph/0211018].
 - [9] T. Becher, R. J. Hill and M. Neubert, arXiv:hep-ph/0308122; T. Becher, R. J. Hill, B. O. Lange and M. Neubert, arXiv:hep-ph/0309227.
 - [10] R. J. Hill, private communication.
 - [11] For a recent discussion on this point see I. Z. Rothstein, arXiv:hep-ph/0308266.
 - [12] M. Beneke and T. Feldmann, arXiv:hep-ph/0311335.